

Treatment-free remission for chronic myeloid leukaemia (CML)

Treatment-free remission (TFR) is when your CML stays under control without the need to keep taking medicines. It might be possible for some people with CML who have a deep, long-term response to tyrosine kinase inhibitors (TKIs). Find out about the risks and benefits, and what happens if you try it.

Summary

Do not stop CML treatment without discussing it with your haematology team.

- Treatment-free remission (TFR) is when your CML stays under control without the need to keep taking medicines.
- It might be an option for people who have been taking TKIs for 3 to 5 years or more and have been in a deep molecular response for at least 2 years.
- If TFR is an option for you, your haematology team should talk to you about the benefits and drawbacks. You should decide together if it's something you'd like to try.
- If you try TFR, you have regular check-ups to monitor your response.
- Around 50 or more in every 100 people who try TFR keep their response to treatment. But up to 50 in every 100 people may lose their response.
- If you lose your response to treatment, restarting TKI treatment is usually very effective.
- Around 20 to 30 in every 100 people who stop TKIs get muscle pain and joint stiffness. This is called treatment discontinuation syndrome. It usually goes away on its own, but some people might need a short course of treatment.

[Download our factsheet about TFR](#) 

[Download our leaflet on TFR](#) 

[Order our TFR leaflet in print](#) 

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

What is treatment-free remission?

Treatment-free remission (TFR) is when your CML stays under control without the need to keep taking medicines. It may be possible for some people with CML who have had a deep, long-lasting response to treatment with tyrosine kinase inhibitors (TKIs).

[Back to top](#)

Who may be able to try TFR?

TFR is not possible for everyone. But it might be an option if:

- Your CML is in the chronic phase and has never been in the blast phase.
- You've been on TKI treatment for at least 3 years (but ideally 5 years)
- You've been in a deep molecular response for at least 2 years.
- You've never had to stop taking a TKI because it didn't work well. (You might still be able to try TFR if you had to stop a TKI because of side effects.)
- You do not have a genetic change in your *BCR-ABL1* gene that makes it resistant to treatment.
- Your hospital has quick access to accurate molecular testing (PCR) to monitor your CML.

[Back to top](#)

Benefits and drawbacks of TFR

If TFR is an option, your haematology team should talk to you about the benefits and drawbacks. You should decide together whether it's something you'd like to try. If you feel anxious and you are not comfortable trying it, let them know. Your haematology team shouldn't force TFR on you if you are not comfortable with it.

Benefits of trying TFR

- ✓ Your CML might stay under control without having to take any medicines. This is the case for around 50 or more in every 100 people who try TFR.
- ✓ You do not have to remember to take medicines every day.
- ✓ You are not at risk of getting any side effects.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- ✓ If you are hoping to get pregnant, TFR might give you the chance to do this safely and stay off TKI treatment during your pregnancy.

Drawbacks of trying TFR

- ✗ Your CML might not stay under control. This happens in up to 50 in every 100 people who try TFR. It usually happens within 6 to 9 months of stopping treatment, although it can happen much later. If your CML does not stay under control, you will need to start treatment again.
- ✗ You have more frequent blood tests and appointments to check if your CML is staying under control.
- ✗ You may get treatment discontinuation syndrome.
- ✗ You may feel anxious about your CML coming back.

[Back to top](#)

What does TFR involve?

If you and your haematology team agree that TFR is right for you, they will explain what the process will involve.

- You might stop your TKI treatment straightaway, or you might take a half-dose for a year before stopping treatment. This may improve the chance of your CML staying in under control once you stop treatment.
- You will have frequent blood tests to check your [molecular response](#). You will also have check-ups to look for treatment discontinuation syndrome.

[Back to top](#)

Treatment discontinuation syndrome

Treatment discontinuation syndrome is muscle pain and stiffness that develops after you stop treatment with TKIs. Around 20 to 30 in every 100 people who stop TKIs get it. Doctors do not know exactly why it happens. It usually starts within days or weeks of stopping your TKI.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Treatment discontinuation syndrome is usually mild and goes away on its own. But some people may need a short course of treatment with paracetamol, non-steroidal anti-inflammatory medicines like ibuprofen or diclofenac, or sometimes steroids (prednisolone).

[Back to top](#)

If CML does not stay under control

If you try TFR, your haematology team will monitor your [molecular response](#) closely. If you lose a major molecular response, you will usually need to start treatment again. This should happen within 4 weeks of losing your response. Most people restart the same TKI they were on before, but your haematologist might suggest a different one.

If you do not achieve a major molecular response within 6 months of restarting treatment, your doctor is likely to check for changes in your *BCR-ABL1* gene. These could affect how well it responds to treatment. They might suggest a different treatment option.

It can be very worrying if your CML does not stay under control, but restarting TKI treatment is usually very effective.

[Back to top](#)

Sources we used to develop this information

Annunziata M, Bonifacio M, Breccia M, Castagnetti F, Gozzini A, Iurlo A, et al. Current Strategies and Future Directions to Achieve Deep Molecular Response and Treatment-Free Remission in Chronic Myeloid Leukemia. *Front Oncol* 2020;10:883.

Baccarani M, Cortes J, Pane F, Niederwieser D, Saglio G, Apperley J, et al. Chronic myeloid leukemia: an update of concepts and management recommendations of European LeukemiaNet. *Journal of clinical oncology*. 2009 Dec 12;27(35):6041.

Bonifacio M, Stagno F, Scaffidi L, Krampera M, Di Raimondo F. Management of chronic myeloid leukaemia in advanced phase. *Front Oncol* 2019;9: Art 1132.

Du Z, Lovly CM. Mechanisms of receptor tyrosine kinase activation in cancer. *Mol Cancer* 2018;17(1):58.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Garcia-Gutierrez V, Hernandez-Boluda JC. Tyrosine kinase inhibitors available for chronic myeloid leukaemia: Efficacy and safety. *Front Oncol* 2019;9:Art 603.

Healey MA, Allendorf DJ, Borate U, Madan A. CNS Involvement in a Patient with Chronic Myeloid Leukemia. *Case Rep Hematol* 2021;2021:8891376.

Hochhaus A, Baccarani M, Silver RT, Schiffer C, Apperley JF, Cervantes F, et al. European LeukemiaNet 2020 recommendations for treating chronic myeloid leukemia. *Leukemia* 2020;34(4):966-984.

Hochhaus A, Saussele S, Rosti G, Mahon FX, Janssen JJWM, Hjorth-Hansen H, et al; ESMO Guidelines Committee. Chronic myeloid leukaemia: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2017;28(suppl_4):iv41-iv51.

Hughes TP, Mauro MJ, Cortes JE, Minami H, Rea D, DeAngelo DJ, et al. Asciminib in chronic myeloid leukaemia after ABL kinase inhibitor failure. *N Engl J Med* 2019; 381:2315-2326.

Incyte Biosciences UK Ltd. Ponatinib 15 mg tablets. Summary of Product Characteristics. July 2022. Available at: <https://www.medicines.org.uk/emc/product/1212/smpc> [Last accessed 9/11/23]

Jabbour E, Kantarjian H, Cortes J. Use of second- and third-generation tyrosine kinase inhibitors in the treatment of chronic myeloid leukaemia: an evolving treatment paradigm. *Clin Lymphoma Myeloma Leuk* 2015;15(6):323-334.

Janssen L, Blijlevens NMA, Drissen MMCM, Bakker EA, Nuijten MAH, Janssen JJWM, et al. Fatigue in chronic myeloid leukemia patients on tyrosine kinase inhibitor therapy: predictors and the relationship with physical activity. *Haematologica* 2021;106(7):1876-1882.

Medac. Hydroxycarbamide 500 mg capsules. Summary of Product Characteristics. February 2023. Available at: <https://www.medicines.org.uk/emc/product/254/smpc> [Last accessed 07/11/2023]

Mylan. Dasatinib 100 mg tablet. Summary of Product Characteristics. 25 July 2023. Available at: <https://www.medicines.org.uk/emc/product/14399/smpc>. [Last accessed: 9/11/23]

National Institute for Health and Care Excellence (NICE). Asciminib for treating chronic myeloid leukaemia after 2 or more tyrosine kinase inhibitors. Technology appraisal

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

guidance [TA813]. Published: 03 August 2022. Available at: <https://www.nice.org.uk/guidance/ta813>

National Institute for Health and Care Excellence (NICE). Bosutinib for previously treated chronic myeloid leukaemia. Technology appraisal guidance [TA401] Published: 24 August 2016. Available at: <https://www.nice.org.uk/guidance/ta401>

National Institute for Health and Care Excellence (NICE). Dasatinib, nilotinib and high-dose imatinib for treating imatinib-resistant or intolerant chronic myeloid leukaemia. Technology appraisal guidance [TA425]. Published: 21 December 2016. Available at: <https://www.nice.org.uk/guidance/ta425>

National Institute for Health and Care Excellence (NICE). Dasatinib, nilotinib and imatinib for untreated chronic myeloid leukaemia. Technology appraisal guidance [TA426] Published: 21 December 2016. Available at: <https://www.nice.org.uk/guidance/ta426>

National Institute for Health and Care Excellence (NICE). Ponatinib for treating chronic myeloid leukaemia and acute lymphoblastic leukaemia. Technology appraisal guidance [TA451] Published: 28 June 2017. Available at: <https://www.nice.org.uk/guidance/ta451> [Last accessed 9/11/23]

Novartis Pharmaceuticals. Imatinib 100 mg tablets. Summary of Product Characteristics. October 2023. Available at: <https://www.medicines.org.uk/emc/product/7779/smpc> [Last accessed 9/11/23]

Novartis Pharmaceuticals. Nilotinib 150 mg capsules. Summary of Product Characteristics. October 2023. Available at: <https://www.medicines.org.uk/emc/product/5852/smpc> [Last accessed 9/11/23]

Pfizer Limited. Bosutinib 100 mg tablet. Summary of Product Characteristics. 15 May 2023. Available at: <https://www.medicines.org.uk/emc/product/3147/smpc>. [Last accessed: 9/11/23]

Radvoyevitch T, Jankovic GM, Tiu RV, Sauntharajah Y, Jackson RC, Hlatky LR, et al. Sex differences in the incidence of chronic myeloid leukemia. *Radiat Environ Biophys* 2014;53(1):55-63.

Saglio G, Gale RP. Prospects for achieving treatment-free remission in chronic myeloid leukaemia. *Br J Haematol* 2020;190(3):318-327.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Smith G, Apperley J, Milojkovic D, Cross NCP, Foroni L, Byrne J, et al; British Society for Haematology. A British Society for Haematology Guideline on the diagnosis and management of chronic myeloid leukaemia. *Br J Haematol* 2020;191(2):171-193. Soverini S, Mancini M, Bavaro L, Cavo M, Martinelli G. Chronic myeloid leukaemia: the paradigm of targeting oncogenic tyrosine kinase signalling and counteracting resistance for successful cancer therapy. *Mol Cancer* 2018;17(1):49

Swerdlow SH, Campo E, Harris NL, Jaffa ES, Pileri SA, Stein H, Thiele J (Eds): WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues (Revised 4th edition). IARC: Lyon 2017.

Westerweel PE, Te Boekhorst PAW, Levin MD, Cornelissen JJ. New approaches and treatment combinations for the management of chronic myeloid leukaemia. *Front Oncol* 2019;9: Art 665.

Need support?

You are not alone. We're here for you whether you have a diagnosis yourself or know someone who has. If you'd like advice, support, or a listening ear, call our freephone helpline on 08088 010 444 or send a WhatsApp message to 07500 068 065.

[Helpline and WhatsApp →](#)

Help us improve our information

We aim to provide information that's reliable, up-to-date, and covers what matters to you. Please complete our short survey to help us improve our information and make sure it meets your needs.

[Complete our short survey →](#)

About our information

This information is aimed at people in the UK. We do our best to make sure it is accurate and up to date but it should not replace advice from your health professional. Find out more [about our information](#).

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Page last reviewed: 31 March 2024

Updated January 2026

Next review due: 31 March 2027



<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/types/chronic-myeloid-leukaemia-cml/treatment-free-remission-for-chronic-myeloid-leukaemia-cml/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)