

Zanubrutinib

Zanubrutinib is a targeted medicine used to treat some people with chronic lymphocytic leukaemia (CLL). We cover what it is, who might have it and how to take it. We also cover its main side effects and any special precautions you need to take while you're on it.

Summary

- Zanubrutinib is a targeted medicine used to treat chronic lymphocytic leukaemia (CLL).
- It comes as capsules that you take every day. The number of capsules you need to take and how often should be printed on the named label on your medicine packaging.
- You will have blood tests to check how well your CLL is responding to zanubrutinib.
- Your haematology team will monitor how your body is coping with treatment.
- Side effects of zanubrutinib are usually mild to moderate. Some people have mild side effects, whereas others get more severe side effects.
- Some side effects can be serious if they are not treated promptly. Contact your haematology team straight away if:
 - You have bleeding such as blood in your poo or pee, nosebleeds, bleeding gums, vomiting or coughing up blood, heavy periods or bruising.
 - You think you might have an infection.
 - You have difficulty breathing, shortness of breath or your breathing is very fast.
 - Your heart rate is fast or uneven, or your pulse is weak or uneven.

Tell your haematology team if you get any other side effects. They may be able to suggest things to help.

[Download our leaflet about zanubrutinib](#) 

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

What is zanubrutinib?

Zanubrutinib is a targeted medicine that blocks a protein called BTK. This protein helps CLL cells to grow and multiply. Blocking it stops CLL cells from growing and dividing, leading to their death.

The brand name for zanubrutinib is Brukinsa.

[Back to top](#)

Who might take it?

Zanubrutinib is suitable for adults. Your haematology team might recommend it for you if:

- You have CLL and have not had treatment before **and**:
- You have a genetic change in your CLL cells called *17p* deletion or *TP53* mutation
or
- Other treatments are unsuitable for you.
- You have had treatment, but your CLL didn't respond to it or came back after treatment. This is called refractory or relapsed CLL.

[Back to top](#)

Before taking zanubrutinib

Before starting zanubrutinib, your haematology team will carry out checks to ensure it is suitable for you. These include:

- Blood tests to check your blood cell counts, kidney function and liver function
- Blood tests to check whether you have any infections that could flare up during treatment, such as viral hepatitis
- Checks of your heart health

Your team may recommend other tests, depending on your circumstances.

Things to tell your haematology team

Your haematology team may ask if there is any possibility you might be pregnant. Zanubrutinib should not be taken during pregnancy.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Some medicines, supplements and food may interact with zanubrutinib. Or they may increase your risk of side effects. It is important to tell your haematology team about any medicines or supplements you are taking. This includes prescribed medicines and medicines you have bought yourself without a prescription. Examples include:

- Herbal medicines such as St John's Wort
- Some blood thinners or medicines used to prevent blood clots, like warfarin
- Some medicines used to treat infections, including some antibiotics, antifungal and antiviral medicines
- Some medicines used to treat a rare sleep problem called narcolepsy
- Some medicines that lower your immune system
- Some medicines used to treat heart conditions and high blood pressure, including high blood pressure in your lungs
- Some medicines used to treat epilepsy
- Some medicines used to treat schizophrenia, severe depression or obsessivecompulsive disorder
- Some medicines used to treat cancer
- Some anti-sickness medicines that are used alongside chemotherapy
- Some opioid medicines that are used for pain relief
- Some medicines used to treat reflux or heartburn

Zanubrutinib capsules contain several ingredients. Tell your haematology team if you have any allergies or sensitivities.

[Back to top](#)

How do you take it?

Zanubrutinib comes as white, oval-shaped capsules that you take by mouth. You take zanubrutinib at home (or anywhere you are staying).

- Swallow your zanubrutinib capsules whole with a glass of water. You can take them with or without food.
- Do not open, break, or chew your capsules and do not dissolve them in water.
- Try to take your capsules at about the same time each day. If you take zanubrutinib twice a day, take it at the same time in the morning and evening.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- Avoid grapefruit or bitter oranges (Seville oranges), including the fruit, fruit juice, or foods or supplements containing them. This is because they affect how much zanubrutinib your body absorbs.
- Do not take supplements containing St John's Wort, fish oil, vitamin E or flaxseed while taking zanubrutinib. This is because they affect how much zanubrutinib your body absorbs.

If you forget to take your capsules

If you forget to take your zanubrutinib:

- Skip the missed dose
- Take your next dose at the usual time

It is important not to take a double dose. If you've taken more zanubrutinib than you should, talk to your doctor, nurse or pharmacist. Or go to the hospital immediately. Take your medicine and the leaflet from its box with you.

[Back to top](#)

Dose

Zanubrutinib comes as capsules containing 80mg of zanubrutinib. The number of capsules you need to take should be printed on the medicine packaging. If you are not sure, ask your doctor or nurse.

Most adults with CLL take 320mg of zanubrutinib a day. You either take it as:

- **Four** capsules **once** a day or
- **Two** capsules **twice** a day. You usually leave around 12 hours between each dose.

You can store zanubrutinib at room temperature. Keep it safe where children cannot see or reach it.

[Back to top](#)

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

How long to take it for

Do not stop taking zanubrutinib without discussing it with your haematology team.

Most people carry on taking zanubrutinib for as long as it is working. Your haematology team may suggest stopping zanubrutinib and trying a different treatment if:

- It is not working well
- It stops working
- You are having side effects that are difficult to cope with
- Check-ups show it is affecting other parts of your body, like your heart, liver or kidneys
- You become pregnant

[Back to top](#)

Monitoring

Your haematology team will check how well zanubrutinib is working and how well your body is coping with it.

- You'll have regular blood tests to check your blood cell counts.
- You'll also have occasional blood tests to check how well your liver and kidneys are working.
- You'll have regular blood pressure checks. Your team may ask you to take your blood pressure at home and keep a record of it.
- Your doctor or nurse will examine you to check for signs and symptoms of:
 - Bleeding.
 - Infection.
 - Heart problems.
 - A rare reaction called tumour lysis syndrome.
 - Changes in your skin.
- If needed, you may have other examinations and scans to check the size of your lymph nodes and organs, like your spleen.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

[Back to top](#)

Precautions

There are some precautions to be aware of when you are taking zanubrutinib.

- If you need surgery or to have a tooth out, you may need to stop zanubrutinib for 3 to 7 days before and after. This is to reduce your risk of bleeding. Your doctor will tell you exactly when to stop taking zanubrutinib and when to start it again.
- Zanubrutinib can make you vulnerable to serious infections. You might need to take medicines to prevent infections while taking zanubrutinib.
- You should avoid eating grapefruit and starfruit (carambola), including the fruit, fruit juice, or foods or supplements containing them.
- You should avoid eating bitter oranges (Seville oranges), including the fruit, bitter orange juice or food made with bitter oranges, like marmalade.
- You should avoid supplements such as fish oil, flaxseed, or vitamin E preparations.
- Zanubrutinib might make you very tired, low in energy, or dizzy. This might affect your ability to drive or operate machinery.
- Other medicines might interact with zanubrutinib. Your haematology team will tell you whether or how to take other medicines while you are on zanubrutinib.
- If you've had viral hepatitis in the past, it might flare up during zanubrutinib treatment. You'll have a blood test before you start treatment. If this shows you've had viral hepatitis, your team will monitor you for flare-ups. You might be referred to a liver specialist for treatment.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)



"If your GP prescribes you a new medicine, ask them to check that it's safe to take with zanubrutinib."

— Helen, living with CLL since 2015

[Back to top](#)

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Fertility, pregnancy and breastfeeding

We do not know what effect zanubrutinib has on fertility or pregnancy. Based on animal studies, it may damage sperm, increase pregnancy loss, and harm unborn babies.

- If you could get pregnant, it's important to use effective contraception while you are taking zanubrutinib and for 1 month after stopping treatment.
- We do not know whether zanubrutinib affects the way hormonal contraceptives work. So, you need to use a barrier method like a condom, diaphragm or cap.
- If you're taking zanubrutinib and you think you might be pregnant, tell your haematology team as soon as possible.
- If you're planning to get pregnant or to make someone pregnant in the future, talk to your haematology team before starting treatment. They will talk to you about your treatment options.

Breastfeeding

We do not know if zanubrutinib passes into breastmilk or what effect it might have on breastfed babies or children.

You should not breastfeed while you are taking zanubrutinib.

[Back to top](#)

Possible side effects

You may get some side effects while you are taking zanubrutinib. Some people have very few side effects, whereas others get more serious side effects. They may differ from day to day.

Your haematology team should tell you what side effects to look out for and who to contact if you need to.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

If you are worried about any symptoms or side effects, contact your haematology team. You do not have to wait until your next appointment.

Your haematology team might:

- Prescribe medicines to help manage some side effects
- Suggest things you can do to help you cope with side effects
- Lower your dose of zanubrutinib if your side effects are serious or difficult to cope with
- Recommend pausing or stopping zanubrutinib, if needed

You can also report side effects via the [MHRA Yellow Card Scheme](#).

[Back to top](#)

Side effects you might notice

This is not a full list of all the side effects that can happen. The patient information leaflet in your medicine package has more information. Or you can find the leaflet in the [Electronic Medicines Compendium](#).

Here, we list some of the more common side effects of zanubrutinib. These affect at least 1 in every 100 people who take zanubrutinib.

- **Infections.** Contact your team straight away if you think you have an infection.

Look out for:

- A high temperature (38°C or higher)
- Shivering, chills or sweating
- Sore throat, sneezing, blocked or runny nose or earache
- Cough, chest pain or shortness of breath
- Burning or stinging when you pee, or peeing more often than usual
- Sickness or diarrhoea
- Cold sores, or ulcers or blisters around your mouth or genitals

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- **Unexpected or unusual bruising or bleeding**, like nosebleeds, bleeding gums or heavy periods. Look out for blood in your poo or pee, or red or purple spots under your skin. This may look different in black or brown skin. Tell your team if you notice any of these.
- **Tummy problems.** You might get diarrhoea, constipation, or tummy ache. You might feel sick or be sick. If you have these side effects, your team can give you medicines to help.
- **Muscle, bone or joint pain.** Talk to your team if this becomes troublesome.
- **Changes to your skin.** You might get a rash. This might look different on black or brown skin.
- **Fatigue.** You might experience extreme tiredness or lack of energy.
- **Dizziness.** You might feel lightheaded.
- **Swelling.** You might have swelling of your feet, ankles or hands.
- **Heart problems.** Contact your doctor if you get a fast or irregular heartbeat or a heart flutter. They will check your heart function.

Protecting yourself from skin cancer

Zanubrutinib can increase your risk of getting some types of skin cancer. It is important to protect your skin from the sun by keeping your skin covered, wearing a wide-brimmed hat, and using a high-factor sunscreen (SPF 30 or higher).

Check for symptoms like a growth, an unusual patch (this can vary in size, colour and texture), a new mole or a change in an existing mole. If you notice any of these, contact your GP.

[Back to top](#)

Side effects your team might notice

There might be some side effects that your haematology team find during regular checks and blood tests. These include:

- Low levels of white blood cells called neutrophils (neutropenia). This makes you more vulnerable to infections. You might need treatment to prevent or treat infections.
- Low levels of red blood cells (anaemia).
- Low levels of platelets (thrombocytopenia). This can cause unusual bruising and bleeding because platelets help your blood to clot. You might need to have a transfusion.
- High blood pressure. If this happens, you might need to take blood pressure medicines.
- High levels of white blood cells. This might happen when you first start zanubrutinib. Your white blood cells may go up before they come down again. It doesn't mean your CLL is getting worse.

[Back to top](#)

Rare side effects to know about

Rarely, zanubrutinib might cause a side effect called tumour lysis syndrome.


This is a serious side effect of some cancer treatments. It happens if a large number of cancer cells die quickly and release chemicals into your bloodstream.

This affects fewer than 1 in every 100 people taking zanubrutinib. So, it does **not** happen to more than 99 in every 100 people. However, if it does happen, it can be serious, so it's sensible to know what to look out for.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)



"If you have to go to A&E, tell the staff you have blood cancer and the treatment you're on. Your team might give you a Patient Alert Card to keep with you. In a medical emergency, these cards let people know what medicine you're taking and help keep you safe."

— *Steve, living with CLL since 2011*

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Contact your doctor immediately if you develop any of the following while taking zanubrutinib:

- Fever or chills
- Diarrhoea, feeling sick or being sick
- Muscle or joint pain
- Cloudy or dark-coloured pee
- Feeling weak, exhausted or confused
- Irregular heartbeat or shortness of breath
- Pain or bloating in your tummy
- Seizures (fits)

[Back to top](#)

If zanubrutinib doesn't work

If zanubrutinib is not working well for you, your doctor will discuss your options with you. They might suggest changing your dose, pausing or stopping treatment.

They will talk to you about other treatment options that might be suitable for you.

[Back to top](#)

Sources we used to develop this information

Beigene UK Limited. Brukinska (zanubrutinib) 80 mg hard capsules - Summary of product characteristics. 05 December 2024. Available at: <https://www.medicines.org.uk/emc/product/14001/smhc>

Beigene UK Limited. Brukinska (zanubrutinib) 80 mg hard capsules. Package leaflet: information for the patient. October 2024. Available at: <https://www.medicines.org.uk/emc/product/14001/pl>

Bennett R, Seymour JF. Update on the management of relapsed/refractory chronic lymphocytic leukemia. Blood Cancer Journal. 2024 Feb 21;14(1):33. <https://doi.org/10.1038/s41408-024-01001-1>

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Molica S, Tam C, Allsup D, Polliack A. Advancements in the treatment of CLL: The rise of zanubrutinib as a preferred therapeutic option. *Cancers*. 2023 Jul 23;15(14):3737. <https://doi.org/10.3390/cancers15143737>

Narang M, Horn C, Lee E. Clinician Perspective on Once-Daily Zanubrutinib Dosing for B-Cell Malignancies at a Single Center. *Clinical Medicine Insights: Oncology*. 2024 Sep;18:11795549241275665. <https://10.1177/11795549241275665>

National Institute for Health and Care Excellence (NICE). Zanubrutinib for treating chronic lymphocytic leukaemia. 2023. Technology appraisal guidance [TA931]. 22 November 2023. Available at: <https://www.nice.org.uk/guidance/ta931>

Tam C, Thompson PA. BTK inhibitors in CLL: second-generation drugs and beyond. *Blood Advances*. 2024 May 14;8(9):2300-9. <https://doi.org/10.1182/bloodadvances.2023012221>

Need support?

You are not alone. We're here for you whether you have a diagnosis yourself or know someone who has. If you'd like advice, support, or a listening ear, call our freephone helpline on 08088 010 444 or send a WhatsApp message to 07500 068 065.

[Helpline and WhatsApp →](#)

Help us improve our information

We aim to provide information that's reliable, up-to-date, and covers what matters to you. Please complete our short survey to help us improve our information and make sure it meets your needs.

[Complete our short survey →](#)

About our information

This information is aimed at people in the UK. We do our best to make sure it is accurate and up to date but it should not replace advice from your health professional. Find out more [about our information](#).

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Page last reviewed: 31 August 2025

Updated February 2026

Next review due: 31 August 2028



<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/zanubrutinib/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)