

Azacitidine tablets

Azacitidine tablets are a type of maintenance therapy used to treat some people with acute myeloid leukaemia (AML).

Summary

- Azacitidine tablets are used as a maintenance treatment for adults with [AML](#).
- Azacitidine is a type of chemotherapy medicine that can turn on cancer-suppressing genes.
- Most people take one tablet a day for 14 days, followed by 14 days off. Your dose should be printed on the named label on the medicine packaging.
- You will have regular blood tests to check how well your AML is responding to azacitidine, and to check your blood counts.
- You may get some side effects while you are taking azacitidine tablets. Some people may have very few side effects, whereas other people experience more serious side effects.
- Some side effects can be serious if they are not treated promptly. Contact your haematology team straight away if you get:
 - A fever of 38°C or higher, shivers, shakes or chills
 - Achy or very tired, feel generally unwell, or poor appetite
 - A cough, chest pain, breathlessness or wheezing
- Tell your haematology team if you get any other side effects. They may be able to suggest things to help.

Do not stop taking azacitidine tablets without talking to your haematology team.

[Download our leaflet about azacitidine tablets](#) 

About azacitidine tablets

Azacitidine is a chemotherapy medicine that can turn on genes that have been turned off. This includes cancer-suppressing genes that are often turned off in AML cells. Azacitidine can also stop AML cells making proteins. This can stop cancer cells growing and dividing.

The brand name for azacitidine tablets is Onureg.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Azacitidine tablets are used as a maintenance treatment for adults with AML. Your haematology team might recommend them for you if:

- Your AML went into complete remission after induction or consolidation therapy, even if your blood counts have not recovered completely **AND**
- You cannot currently have, or do not want, a [stem cell transplant](#)

[Back to top](#)

Before taking azacitidine tablets

Before starting your course of azacitidine, your haematology team will carry out checks to make sure it's suitable for you. These usually include:

- Blood tests to measure your blood counts
- Blood tests to check your liver and kidney function
- Bone marrow tests to check that your AML is still in remission

[Back to top](#)

Taking azacitidine tablets

Azacitidine tablets come as film-coated tablets containing 200mg or 300mg of azacitidine. Most people take one tablet a day for 14 days of each month. Your dose should be printed on the named label on the medicine packaging. If you are not sure, ask your doctor or nurse.

- Your doctor will also give you tablets to help stop you feeling sick or being sick. You should take these 30 minutes before you take your azacitidine tablets. Usually, you only need to take the anti-sickness tablets for your first two treatment cycles, but your doctor will let you know if you need to take them for longer.
- You should take azacitidine as prescribed so it can work as well as possible. Try and take it at the same time of day if you can.
- Swallow the azacitidine tablet whole with a full glass of water. To make sure you get the right dose, do not break, crush, dissolve or chew the tablet.
- You can take the tablet with food or in between meals, whichever is best for you.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- If you are sick after taking a tablet, do not take another dose on the same day. Instead, wait until the next day and take your dose as normal. It is important not to take a double dose.
- If you forget to take your azacitidine tablet, take it as soon as you remember on the same day. If it is the next day, just take your usual dose at the usual time. Do not take a double dose to make up for a forgotten tablet.
- If powder from a broken tablet touches your skin, wash it straight away and thoroughly with soap and water. If the powder gets into your eyes, nose or mouth, flush the area thoroughly with water.

Dose

Most adults with AML take 300mg azacitidine tablets. You usually have them in a treatment cycle of 28 days.

- You take azacitidine tablets every day for the first 14 days.
- This is followed by 14 days where you do not take the azacitidine tablets.

Your haematology team may suggest a different dose, depending on your circumstances. They will tell you what dose they recommend for you.

[Back to top](#)

Precautions

There are several precautions to be aware of when you are taking azacitidine tablets.

- Azacitidine tablets may make you feel tired or weak or make it difficult to concentrate. Take care if you are driving or using tools or machinery.
- Azacitidine tablets contain lactose. If you think you may have lactose intolerance, let your haematology team know.
- Tell your haematology team about any medicines or herbal supplements that you are taking.

[Back to top](#)

Pregnancy, breastfeeding and fertility

Azacitidine tablets may harm unborn babies, especially during the early stages of pregnancy.

- If you could get pregnant, it is important to use effective contraception while you are taking azacitidine tablets and for 6 months after you stop.
- If you could make someone pregnant, it is important to use effective contraception while you are taking azacitidine tablets and for 3 months after you stop.
- If you are taking azacitidine tablets and think you may be pregnant, tell your haematology team as soon as possible. They may recommend stopping treatment for a while or switching to a different treatment.
- If you are planning to get pregnant, or make someone pregnant, tell your haematology team. They can discuss your treatment options with you.

Breastfeeding

Scientists are not sure if azacitidine passes into breastmilk. If it does, it could be a risk for breastfed babies or children. You should not breastfeed when taking azacitidine.

[Back to top](#)

Monitoring

During your treatment with azacitidine tablets, you will have blood tests every 2 to 4 weeks.

Your haematology team will use the results of these tests to:

- Check how well azacitidine tablets are working for you
- Make sure your blood counts have not dropped too low

[Back to top](#)

How long to take azacitidine tablets

You usually carry on taking azacitidine tablets unless:

- It stops working well

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- You are having side effects that are difficult to cope with

If this happens, your haematology team might stop azacitidine, adjust your dose, or recommend a different treatment.

Do not stop taking azacitidine tablets without talking to your haematology team.

[Back to top](#)

Side effects

You may get some side effects while you are taking azacitidine tablets. Some people may have very few side effects, whereas other people experience more serious side effects.

Tell your haematology team if you get any side effects. They may be able to suggest things to help.

If you are getting side effects that are difficult to cope with, they might suggest:

- Temporarily stopping treatment
- Lowering your dose of azacitidine
- Changing your treatment so you only take azacitidine tablets for 7 days each cycle

If you are worried about any side effects or symptoms you have, contact your haematology team

[Back to top](#)

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Serious side effects

The following side effects may be serious and require urgent treatment. Contact your doctor or nurse straight away if you have any of these side effects.

- A high temperature (higher than 38°C) with a low white blood cell count (febrile neutropenia). You might also feel shivery or generally unwell.
- Pneumonia (a lung infection). You may get symptoms like:
 - A cough, chest pain
 - A high temperature
 - Shortness of breath, wheezing when you breathe
 - An aching body, feeling very tired
 - Loss of appetite

Azacitidine can lower your blood cell counts. This can mean you may be more likely to get infections, fever, bruising or bleeding. If you get any of these symptoms, tell your doctor or nurse straight away.

[Back to top](#)

Very common side effects

Very common side effects affect more than 10 in every 100 people who are taking azacitidine tablets.

Very common side effects include:

- Feeling sick or being sick
- Diarrhoea
- A low white blood cell count, which might mean that you get infections more often and any infections you get might last longer than usual

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- Feeling tired or weak
- Constipation
- A low platelet count, which might lead to unexpected bruising or bleeding like nosebleeds, bleeding gums, blood spots or rashes
- Pain in your tummy
- Throat, nose or sinus infections
- Stiff or painful joints
- Loss of appetite
- Back pain
- Pains in your arms or legs
- Lung infections

[Back to top](#)

Common side effects

Common side effects affect more than 1 but less than 10 out of every 100 people who are taking azacitidine tablets.

Common side effects include:

- Flu
- Urine infections
- You may have darker or smellier pee, stinging when going for a pee or needing to pee more than usual
- Runny nose or cough
- Anxiety
- Weight loss

[Back to top](#)

If azacitidine tablets do not work

If azacitidine tablets are not working well, your haematology team might suggest increasing your treatment so you take the azacitidine tablets for more than 14 days during each cycle.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

If this is not suitable for you, or does not work, your team will talk to you about your treatment options.

[Back to top](#)

Sources we used to develop this information

Bristol Myers Squibb Pharmaceuticals. Onureg 200 mg film-coated tablets – Patient Information Leaflet [Internet]. 2022. Available from <https://www.medicines.org.uk/emc/product/14217/pil> [Accessed Aug 2024].

Bristol Myers Squibb Pharmaceuticals. Onureg 200 mg film-coated tablets – Summary of Product Characteristics [Internet]. 2021. Available from <https://www.medicines.org.uk/emc/product/14217/smhc> [Accessed Aug 2024].

Bristol Myers Squibb Pharmaceuticals. Onureg 300 mg film-coated tablets – Patient Information Leaflet [Internet]. 2022. Available from <https://www.medicines.org.uk/emc/product/14218/pil> [Accessed Aug 2024].

Bristol Myers Squibb Pharmaceuticals. Onureg 300 mg film-coated tablets – Summary of Product Characteristics [Internet]. 2021. Available from <https://www.medicines.org.uk/emc/product/14218/smhc> [Accessed Aug 2024].

Need support?

You are not alone. We're here for you whether you have a diagnosis yourself or know someone who has. If you'd like advice, support, or a listening ear, call our freephone helpline on 08088 010 444 or send a WhatsApp message to 07500 068 065.

[Talk to us →](#)

Help us improve our information

We aim to provide information that's reliable, up-to-date, and covers what matters to you. Please complete our short survey to help us improve our information and make sure it meets your needs.

[Complete our short survey →](#)

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

About our information

This information is aimed at people in the UK. We do our best to make sure it is accurate and up to date but it should not replace advice from your health professional. Find out more [about our information](#).

Page last reviewed: 30 November 2024

Updated February 2026

Next review due: 30 November 2027

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/azacitidine-tablets/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)