

Venetoclax + obinutuzumab

Venetoclax and obinutuzumab is a combination of medicines used to treat some people with chronic lymphocytic leukaemia (CLL).

Summary

- Venetoclax and obinutuzumab is a combination of medicines. It is used to treat some people with chronic lymphocytic leukaemia (CLL).
- Venetoclax is a targeted treatment. It comes as film-coated tablets that you take by mouth every day.
- Obinutuzumab is an antibody therapy. You have it through a drip into a vein at a hospital day unit.
- You have venetoclax and obinutuzumab treatment in cycles. One cycle is 28 days.
- Your haematology team will monitor how your body is coping with treatment.
- You will also have blood tests to check how well your CLL is responding.
- Some people get side effects. Some people have mild side effects, whereas others get more severe side effects.
- Some side effects can be serious if they are not treated promptly. Contact your haematology team straight away if:
 - You think you might have an infection
 - Your pee is dark or cloudy
 - You have a seizure (fit)
 - You feel weak, exhausted or confused
 - You have pain or bloating in your tummy, diarrhoea or sickness
 - You have chest pain, difficulty breathing or shortness of breath
 - Your heart rate is fast or uneven, or your pulse is weak or uneven
 - You develop a rash or blue, grey or pale, blotchy skin, lips or tongue (on black or brown skin, look on the palms of the hands or soles of the feet)
- Tell your haematology team if you get any other side effects. They may be able to suggest things to help.

[Download our leaflet about venetoclax and obinutuzumab](#) 

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

What is venetoclax and obinutuzumab?

Venetoclax and obinutuzumab is a combination of medicines. It is used to treat some people with [chronic lymphocytic leukaemia \(CLL\)](#).

Venetoclax

Venetoclax is a targeted medicine that helps slow down the progression of CLL. It blocks a protein in CLL cells called BCL-2. This protein stops CLL cells dying when they should. Blocking it makes the cells self-destruct and die.

The brand name for venetoclax is Venclyxto.

Obinutuzumab

Obinutuzumab is an antibody therapy made with antibodies produced in a lab. They are monoclonal antibodies, which means they are all of one type. The antibodies stick to a protein on the surface of CLL cells called CD20. This triggers your immune system to destroy the CLL cells.

The brand name for obinutuzumab is Gazyvaro.

[Back to top](#)

Who might have it?

Venetoclax and obinutuzumab is suitable for adults. Your haematology team might recommend it if you have CLL that needs treatment, and you have not had treatment for it before.

Treatment aims to reduce your symptoms and keep your CLL under control for as long as possible with as few side effects as possible.

[Back to top](#)

Before starting this treatment

Before prescribing venetoclax and obinutuzumab, your haematology team will make sure it is suitable for you.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

You may have:

- Blood tests to check your blood cell counts
- Blood tests to check your liver and kidney function, and the levels of salts and minerals in your body
- Blood tests to check whether you have an active infection or infections that could flare up during treatment, such as viral hepatitis
- Checks of your heart health
- A CT scan or MRI scan to find out if CLL is affecting your organs, like your spleen

Your team may recommend other tests, depending on your circumstances.

Your haematology team may ask if there is any possibility you might be pregnant.

Venetoclax and obinutuzumab is not recommended during pregnancy.

Things to tell your haematology team

Some medicines, supplements and food may affect how well your treatment works. Or they may increase your risk of side effects. It is important to tell your haematology team about any medicines you are taking. This includes prescribed medicines and medicines you have bought yourself without a prescription.

Examples include:

- Herbal medicines such as St John's Wort.
- Some blood thinners or medicines used to prevent blood clots, like warfarin.
- Some medicines to treat infections, including some antibiotics, antifungal and antiviral medicines.
- Some medicines used to treat high blood pressure or heart problems.
- Some medicines used to prevent or treat stroke, or blood clots in your legs or lungs.
- Some medicines used to lower cholesterol.
- Some medicines used to treat epilepsy.
- Some medicines used to treat a rare sleep problem called narcolepsy.
- Blood pressure medicines. You may need to stop these on the day of each obinutuzumab dose, to reduce the chance of low blood pressure.

Venetoclax and obinutuzumab contain several ingredients. Tell your haematology team if you have any allergies or sensitivities.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

[Back to top](#)

How do you have venetoclax?

Venetoclax comes as tablets you take by mouth



- Swallow your tablets whole with a glass of water.
- Do not chew, crush, or break your tablets and do not dissolve them in water.
- Try to take your tablets at about the same time each day. Take them with a meal because this helps your body absorb the right amount of venetoclax.
- You should drink plenty of water to help reduce the side effects of venetoclax. Especially on the days before and the day you start taking it. Also, whenever you increase your dose.
- You need to take medicine to help control your uric acid levels. You'll start this a few days before you begin your course of venetoclax. You carry on taking it until you are on a stable dose of venetoclax, or longer if needed. Your doctor will prescribe it for you.
- Avoid grapefruit, bitter oranges (Seville oranges) or starfruit (carambola) while taking venetoclax. This includes the fruit, fruit juice, or foods or supplements containing them. These foods affect how much medicine your body absorbs.
- Do not take supplements containing St John's Wort while taking venetoclax. This is because it affects how much medicine your body absorbs.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

If you miss a dose

If you forget to take venetoclax:

- Take it as soon as you remember, if it's **within 8 hours** of your dose time.
- If it's **longer than 8 hours** by the time you realise you forgot to take your tablet:
 - Skip the missed tablets
 - Take your next tablets at the usual time the following day
- If you are sick (vomit) after you've taken your tablets, don't take any more tablets that day. Take your next dose at the usual time the following day.

It is important not to take a double dose. If you've taken more venetoclax than you should, talk to your doctor, nurse or pharmacist. Or go to the hospital immediately. Take your medicine and the leaflet from its box with you.

[Back to top](#)

How do you have obinutuzumab?

Obinutuzumab is an antibody infusion. You have it through a drip into a vein.



<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

The day before having obinutuzumab, you may start medicine to control your uric acid levels. This helps prevent a serious side effect called [tumour lysis syndrome](#). Your doctor will prescribe it for you and tell you when to take it.

- You also need to drink plenty of fluids before you have obinutuzumab. This helps prevent serious side effects.
- About an hour before each dose of obinutuzumab, you'll have medicine to help prevent any reactions to it. This may include:
 - Paracetamol.
 - Antihistamines, which may make you feel sleepy or drowsy. You should not drive if you feel this way.
 - Steroids (you might only have these during your first two doses).
- You then have obinutuzumab through a thin tube (cannula). This is usually in your arm or the back of your hand. It usually takes around 4 to 5 hours the first time you have the drip.

If you miss an appointment to have obinutuzumab, your haematology team should rearrange it as soon as possible.

[Back to top](#)

Dose

You start obinutuzumab treatment first. You start with a low dose. You usually start venetoclax 3 weeks after your first dose of obinutuzumab.

You have venetoclax and obinutuzumab treatment in 28-day cycles:

- You have six cycles of venetoclax and obinutuzumab
- You have another six cycles of venetoclax alone

Most people have their dose of venetoclax and obinutuzumab increased over the first two 28-day cycles. Then they have four more cycles of obinutuzumab and venetoclax, followed by six cycles of venetoclax on its own.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Cycle 1

- Day 1: You have 100mg of obinutuzumab very slowly to check for reactions to the drip.
- Day 1 or 2: You have 900mg of obinutuzumab.
- Day 8: You have 1,000mg of obinutuzumab.
- Day 15: You have 1,000mg of obinutuzumab.
- Day 22 to 28: Take two 10mg venetoclax tablets once a day.

Cycle 2

- Day 1 to 7: You have 1,000mg of obinutuzumab on day 1 and take one 50mg venetoclax tablet once a day.
- Day 8 to 14: Take one 100mg venetoclax tablet once a day.
- Day 15 to 21: Take two 100mg venetoclax tablets once a day.
- Day 22 onwards: Take four 100mg venetoclax tablets once a day.

Cycles 3 to 6

- Day 1: You have 1,000mg of obinutuzumab.
- Continue taking four 100 mg venetoclax tablets once a day.

Cycles 7 to 12

Continue taking four 100mg venetoclax tablets once a day.

Obinutuzumab

You start with a low dose infusion. Your doctor usually increases the dose over the first week until you reach the full dose of 1,000mg. Then you have 1,000mg once a month.

Venetoclax

Venetoclax comes as film-coated tablets containing 10mg, 50mg or 100mg of venetoclax. The number of tablets you need to take should be printed on the medicine packaging. If you are not sure, ask your doctor or nurse.

You take venetoclax once a day, starting on a low dose of 20mg. Your doctor usually increases the dose over 4 weeks until you reach the full dose of 400mg. Your dose is increased slowly to help prevent tumour lysis syndrome. You might hear people call this the 'ramping up' stage.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

For the first 4 weeks of venetoclax treatment, you'll get a new pack each week. They usually come as blister packs with the daily dose you need contained within each blister. This will help you take the right dose each day while your doctor is increasing your dose.

[Back to top](#)

Where to have this treatment

You usually have **obinutuzumab** at a hospital day unit and go home the same day. Some people may need a short hospital stay.

You usually go to a hospital day unit for your first dose of venetoclax, and when your dose is increased. This is so you can be monitored closely. Some people may need a short hospital stay.

The rest of the time, you can take **venetoclax** at home (or wherever you are staying). You can store venetoclax at room temperature. Keep your tablets safe where children cannot see or reach them.

[Back to top](#)

How often to have it

In the first month, you have four doses of obinutuzumab. After this, you have an obinutuzumab infusion on day 1 of the next 5 cycles.

During the first few weeks of treatment with venetoclax, you should take it in the morning, with food. This will help with follow-up blood tests while your doctor increases your dose.

From cycle 1, day 22 onwards, you also take venetoclax tablets once a day, every day. Try to take your tablets at the same time every day, with food.

[Back to top](#)

How long to have it for

Do not stop taking venetoclax without discussing it with your haematology team.

Treatment usually lasts for 1 year. You only have obinutuzumab for the first 6 months of this.

Once you have completed your treatment, you may go on to [active monitoring](#). If you need treatment again in the future, you may be offered the same treatment again if it worked well for you.

Your haematology team may suggest stopping venetoclax and obinutuzumab early and trying a different treatment if:

- It is not working well
- It stops working
- You are having side effects that are difficult to cope with
- Check-ups show it is affecting other parts of your body, like your heart or kidneys
- You become pregnant

[Back to top](#)

Monitoring

During each dose of **obinutuzumab**, a nurse will regularly check your blood pressure, temperature and oxygen levels.

When you start **venetoclax** and for each dose increase, you'll have blood tests:

- Before you take your tablets.
- 6 to 8 hours after taking your tablets. Or more often, if your team think you need it.
- Around 24 hours after taking your tablets. You usually come back to the hospital for this.

The exact timing of your blood tests may vary. It may depend on where you live and how long it takes you to travel to and from the clinic.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Once you are on a stable dose of **venetoclax and obinutuzumab**, your haematology team will continue to monitor you. This is to check how well your treatment is working and how well your body is coping with it.

- You'll have regular blood tests to check:
 - Your blood cell counts.
 - How well your liver and kidneys are working.
 - The levels of salts, minerals and uric acid in your body.
- You'll have regular blood pressure checks.
- Your doctor or nurse will examine you to check for signs of infection, or signs of other side effects.
- You may have other examinations and scans to check the size of your lymph nodes and organs, like your spleen.
- Your team might recommend other tests, depending on your circumstances.

[Back to top](#)

Precautions

There are some precautions to be aware of when you are taking venetoclax and obinutuzumab.

- Both venetoclax and obinutuzumab can make you vulnerable to infections. You might need to take medicines to prevent infections while you're having treatment.
- You should avoid eating grapefruit and starfruit (carambola). This includes the fruit, fruit juice, or foods or supplements containing them.
- You should avoid eating bitter oranges (Seville oranges). This includes the fruit, bitter orange juice or food made with bitter oranges, like marmalade.
- Venetoclax may make you very tired or dizzy. This might affect your ability to drive or operate machinery.
- Obinutuzumab can lower your blood pressure. If you are on blood pressure medication, your haematology team may tell you not to take it the day before and the day of obinutuzumab treatment.
- Other medicines might interact with venetoclax and affect how well it works. Your haematology team will tell you whether or how to take other medicines while you're taking it. Or whether you need a different dose of venetoclax than usual.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- You should not have live vaccines when you have CLL or when you are on venetoclax and obinutuzumab. It is safe to have non-live vaccines. But they might not work as well as they do in other people.



"If you're on venetoclax and obinutuzumab and your GP prescribes you a new medicine, ask them to check that it's safe to take while on your treatment. Your pharmacist can check for interactions with your other medications, too."

— *Helen, living with CLL since 2015*

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Protecting yourself from skin cancer

Venetoclax and obinutuzumab might increase your risk of getting some types of skin cancer.

It is important to protect your skin from the sun by keeping your skin covered, wearing a wide-brimmed hat, and using a high factor suncream (SPF 30 or higher).

Check for symptoms like a growth, an unusual patch (this can vary in size, colour and texture), a new mole or a change in an existing mole. If you notice any, contact your GP.

Precautions to prevent tumour lysis syndrome

When you're taking venetoclax and obinutuzumab, you'll have medicine to help prevent a serious side effect called [tumour lysis syndrome](#). You need to drink plenty of water too, as this will also help prevent it. Try to aim for 1.5 to 2 litres of fluid every day. Especially:

- For 2 days before starting venetoclax
- The day you start venetoclax
- Every time you increase your dose
- Before you have each dose of obinutuzumab

You can include non-alcoholic drinks and non-caffeinated drinks in this amount. But not grapefruit, Seville orange or starfruit juice.

If you're struggling to drink enough, your haematology team might give you fluid through a drip before your treatment.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)



"The weeks when venetoclax is introduced and ramped up are extra busy. You need to have blood tests at certain times before and after each new venetoclax dose to check you aren't developing tumour lysis syndrome."

— Jane, living with CLL since 2004

[Back to top](#)

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Fertility, pregnancy and breastfeeding

We do not know what effect venetoclax and obinutuzumab has on pregnancy. Based on animal studies, venetoclax may damage sperm and harm unborn babies. Also based on animal studies, obinutuzumab may cause very low white blood cell counts in babies who were exposed to it during pregnancy.

- If you could get pregnant, it's important to use effective contraception:
 - While you are taking venetoclax and for at least 30 days after stopping treatment.
 - While you are having obinutuzumab and for 18 months afterwards.
- We do not know whether venetoclax affects the way hormonal contraceptives work. So, you need to use a barrier method like condoms, a diaphragm or cap when you're on venetoclax and obinutuzumab.
- If you're on venetoclax and obinutuzumab and you think you might be pregnant, tell your haematology team as soon as possible.
- Venetoclax may cause male infertility. If you are planning to make someone pregnant in the future, tell your haematology team. They will talk to you about options to preserve your fertility, such as freezing your sperm.
- If you have obinutuzumab when you are pregnant, your baby may have a very low white blood cell count at birth and for months afterwards. Your baby should not have any live vaccines until their blood cell count has recovered.

Breastfeeding

We do not know if venetoclax passes into breastmilk or what effect it might have on breastfed babies or children.

Obinutuzumab passes into breastmilk. We do not know how this might affect breastfed babies or children.

You should not breastfeed:

- While you are on venetoclax
- While you are on obinutuzumab and for 18 months after stopping it

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

[Back to top](#)

Possible side effects

You may get some side effects while you are taking venetoclax and obinutuzumab. Some people have very few side effects, whereas others get more serious side effects. They may differ from day to day.

Your haematology team should tell you what side effects to look out for and who to contact if you need to.

If you are worried about any symptoms or side effects, contact your haematology team. You do not have to wait until your next appointment.

Your haematology team might:

- Prescribe medicines to help manage some side effects
- Suggest things you can do to help you cope with side effects
- Lower your dose of venetoclax, if your side effects are serious or difficult to cope with
- Pause or slow down your obinutuzumab drip, if you have a reaction to it
- Recommend pausing or stopping venetoclax and obinutuzumab, if needed

You can also report side effects via the [MHRA Yellow Card Scheme](#).

[Back to top](#)

Side effects you might notice

This is not a full list of all the side effects that can happen. The patient information leaflets in your medicine packages have more information. Or you can find the leaflets by searching for 'venetoclax' and 'obinutuzumab' at the [Electronic Medicines Compendium](#).

Here, we list some of the more common side effects of venetoclax and obinutuzumab. These affect at least 10 in every 100 people who take venetoclax and obinutuzumab.

Infections

Contact your team straight away if you think you have an infection. Look out for:

- A high temperature (38°C or higher)
 - Shivering, chills or sweating
 - Sore throat, sneezing, blocked or runny nose or earache
 - Cough, chest pain or shortness of breath
 - Burning or stinging when you pee, or peeing more often than usual
- Tummy problems You might get diarrhoea or constipation. You might feel sick or be sick. If you have these side effects, your team can give you medicines to help.

Fatigue, headache or fever

You might experience extreme tiredness or lack of energy. You might get a headache or fever.



"Some people have nausea as a side effect and find that taking venetoclax at bedtime helps, after the ramping up stage ends. It reduces side effects, as they sleep through them. Once I was on 400mg a day, with the agreement of my consultant, I changed to taking my tablets at bedtime."

— *Jane, living with CLL since 2004*

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Reaction to the drip

Some people develop a reaction to the obinutuzumab drip. This can sometimes be serious. You may have symptoms like:

- Breathlessness
- Wheezing, coughing or tightness of the chest
- Rapid heart rate, restlessness and confusion

Tell the day unit team immediately if you develop any symptoms while you are having your rituximab drip.

It's most likely to happen the first time you have obinutuzumab. This is why you have a lower dose at first. You also have your first two doses more slowly than later doses.

Your haematology team will monitor you closely for any signs of a reaction. If needed, they will pause the drip to give you time to recover. They may restart it at a lower rate once you're feeling better. If it is very severe, or it happens again, you may have to stop obinutuzumab.

[Back to top](#)

Side effects your team might notice

There might be some side effects that your haematology team find during regular checks and blood tests. These include:

- Low levels of white blood cells called neutrophils (neutropenia). This can make you more vulnerable to infection.
 - Low levels of red blood cells (anaemia).
 - Low levels of platelets (thrombocytopenia).
 - Changes to the level of salts in your blood, which may be a sign of tumour lysis syndrome.
-

Serious side effects to know about

This is not a full list of all the side effects that can happen. The patient information leaflets in your medicine packages have more information. Or you can find the leaflets by searching for 'venetoclax' and 'obinutuzumab' at the [Electronic Medicines Compendium](#).

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Tumour lysis syndrome

Venetoclax and obinutuzumab is very good at killing cancer cells. It can cause many cancer cells to break down all at once. When this happens, the dying cells release a lot of chemicals into your blood. This can cause a serious condition called tumour lysis syndrome.

Tumour lysis syndrome affects less than 2 in every 100 people taking venetoclax and obinutuzumab. More than 98 in every 100 people do not get it.

Tumour lysis syndrome can make you very unwell while your body tries to cope with changes to your body salts and levels of uric acid:

- Your kidneys may work less well
- Your heartbeat may become irregular
- You may have seizures (fits)


It can happen to anyone having venetoclax and obinutuzumab to treat CLL. But it's more likely to happen if you have lots of CLL cells in your body. It's also more likely if you have other health problems like kidney disease or a swollen spleen. This is because it's harder for your body to cope with all the chemicals from the dead CLL cells.

Starting obinutuzumab before you start venetoclax helps lower your CLL cell number and your chance of tumour lysis syndrome. You are also closely monitored by your haematology team for tumour lysis syndrome. Most people who get tumour lysis syndrome only have changes on blood tests and do not get symptoms. This is known as 'chemical tumour lysis syndrome'. However, you still need to know what to look out for, just in case.

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)



"If you have to go to A&E, tell the staff you have blood cancer and the treatment you're on. Your team might give you a Patient Alert Card to keep with you. In a medical emergency, these cards let people know what medicine you're taking and help keep you safe. "

— Steve, living with CLL since 2011

Contact your haematology team immediately if you develop any of the following symptoms of tumour lysis syndrome:

- **Fever or chills**
- **Diarrhoea, feeling sick or being sick**
- **Muscle or joint pain**
- **Cloudy or dark-coloured pee**

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

- **Feeling weak, exhausted or confused**
- **Irregular heartbeat or shortness of breath**
- **Pain or bloating in your tummy**
- **Seizures (fits)**

The Electronic Medicines Compendium has a [Venetoclax Patient Alert Card](#).

Sepsis

Sepsis is a severe response to an infection. It is a serious condition and needs prompt treatment.

Contact your doctor straight away if you get symptoms of sepsis like:

- **Confusion, slurred speech or not making sense**
- **Blue, grey or pale, blotchy skin, lips or tongue (on black or brown skin look on the palms of the hands or soles of the feet)**
- **A meningitis-like rash, which does not fade when you roll a glass over it**
- **Breathing difficulties, such as breathlessness or rapid breathing**

Heart problems

Obinutuzumab can sometimes cause heart problems. This is more common in people who already have a heart condition. Your team will monitor you for this. Tell them straight away if you have any palpitations, chest pain or breathlessness.

[Back to top](#)

If this treatment doesn't work

If venetoclax and obinutuzumab is not working well for you, your doctor will discuss your options with you. They might suggest changing your dose, pausing or stopping treatment. They will talk to you about other treatment options that might be suitable.

[Back to top](#)

Sources we used to develop this information

AbbVie Ltd. Venclyxto (venetoclax) 10 mg film-coated tablets – Summary of product characteristics. 20 February 2025. Available at: <https://www.medicines.org.uk/emc/product/2267/smpc>

AbbVie Ltd. Venclyxto (venetoclax) 100 mg film-coated tablets – Summary of product characteristics. 26 February 2025. Available at: <https://www.medicines.org.uk/emc/product/10476/smpc>

AbbVie Ltd. Venclyxto (venetoclax) 50 mg film-coated tablets – Summary of product characteristics. 26 February 2025. Available at: <https://www.medicines.org.uk/emc/product/10475/smpc>

AbbVie Ltd. Venclyxto (venetoclax): 10 mg, 50mg, 100mg film-coated tablets. Package leaflet: information for the patient. January 2025. Available at: <https://www.medicines.org.uk/emc/product/2267/pil>

Fischer K, Al-Sawaf O, Bahlo J, Fink AM, Tandon M, Dixon M, Robrecht S, Warburton S, Humphrey K, Samoylova O, Liberati AM. Venetoclax and obinutuzumab in patients with CLL and coexisting conditions. *New England Journal of Medicine*. 2019 Jun 6;380(23):2225-36. Available at: <https://doi.org/full/10.1056/NEJMoa1815281>

Flamingo Pharma (UK) Ltd. Diphenhydramine Hydrochloride 50 mg tablets – Summary of product characteristics. 29 Apr 2022. Available at: <https://www.medicines.org.uk/emc/product/13533/smpc>

National Institute for Health and Care Excellence (NICE). Venetoclax with obinutuzumab for untreated chronic lymphocytic leukaemia. 2020. Technology appraisal guidance [TA663]. 27 February 2019. Available at: <https://www.nice.org.uk/guidance/ta663>

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

Roche Products Limited. Gazyvara 1,000mg concentrate for solution for infusion – Summary of product characteristics. 11 April 2025. Available at: <https://www.medicines.org.uk/emc/product/3279/smpc>

Scottish Medicines Consortium (SMC). Venetoclax (Venclyxto) [SMC 2293]. 6 November 2020. Available at: <https://scottishmedicines.org.uk/medicines-advice/venetoclax-venclyxto-full-smc2293/>

Scottish Medicines Consortium (SMC). Venetoclax (Venclyxto) [SMC 2427]. 8 April 2022. Available at: <https://scottishmedicines.org.uk/medicines-advice/venetoclax-venclyxto-full-smc2427/>

Sportoletti P, Laurenti L, Ferrant E, Montero LF, Mulligan SP, Harrup R, Opat S, Ibatici A, Marasca R, Lefebure M, Boyer M. Reduction of Tumor Lysis Syndrome Risk after Debulking with Obinutuzumab in Patients Treated with Venetoclax-Obinutuzumab in the Randomized Phase 3 CRISTALLO Trial. *Blood*. 2024 Nov 5;144:1873. <https://doi.org/10.1182/blood-2024-198372>

Walewska R, Parry-Jones N, Eyre TA, Follows G, Martinez-Calle N, McCarthy H, Parry H, Patten PE, Riches JC, Hillmen P, Schuh AH. Guideline for the treatment of chronic lymphocytic leukaemia. *Br J Haematol*. 2022 Jun 1;197(5):544-7. <https://doi.org/10.1111/bjh.18075>

Need support?

You are not alone. We're here for you whether you have a diagnosis yourself or know someone who has. If you'd like advice, support, or a listening ear, call our freephone helpline on 08088 010 444 or send a WhatsApp message to 07500 068 065.

[Talk to us →](#)

Help us improve our information

We aim to provide information that's reliable, up-to-date, and covers what matters to you. Please complete our short survey to help us improve our information and make sure it meets your needs.

[Complete our short survey →](#)

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)

About our information

This information is aimed at people in the UK. We do our best to make sure it is accurate and up to date but it should not replace advice from your health professional. Find out more [about our information](#).

Page last reviewed: 31 August 2025

Updated February 2026

Next review due: 31 August 2028

<https://lcdemo-stage.gb.aldryn.io/about-leukaemia/treatments/venetoclax-and-obinutuzumab/>

Leukaemia Care Registered Charity Number 1183890. Scotland Registered Charity Number SC049802

Helpline: [08088 010 444](tel:08088010444)